

ICA International Chemical PRODUCT

1. Virukill™

1) Composition

Didecyldimethyl Ammonium Chloride 120 g/l

2) Properties

Broad spectrum Disinfectant that can be used in manual way, dipping, spray/fogging. Virukill effective in wide range of pH, high and low temperature, and non corrosive.

3) Indications

To disinfect cage, warehouse, squeezing place, equipment, hatchery machine, dan other place in husbandry. Also can be used to disinfect eggs, or mix in drinking water.

4) Dosage and Application :

Needs	Diseases agent	Dosage	How to use
Washing and cleaning the surface	Virus, bacterial, fungi, mycoplasma, yeast dan alga	200 ml per 100 lt of water	Spray 0.5 – 1 L per m ² surface and wash after 30 minutes
disinfect all animal place (cage, dog stable, horse stable, warehouse, squeezing place, laboratory, etc) all surface, equipment, hatchery machine, and rinse water channel		1 lt per 100 lt of water	Pour surface (300-400 ml virukill per m ²) and let it dry
For soak feet and vehicle		1 lt per 25 lt of water	Throw solution if it's dirty
In Drinking water		100 ml per 1000 lt of water	Can be used continuesly
disinfect place before working (related) to animal		1 lt per 100 lt of water	Use in Spray or fogging
disinfect place before working (related) to animal		1 lt per 100 lt of water, 1 – 3 times per day	Use in Spray or fogging
Disinfect eggs.		1 lt per 100 lt of water	Is recommended that solution has the same temperatur with eggs

5) Packaging : 1 box @ 12 x 1 litre, 200 litre